

Jurnal Ilmiah
FARMASI

(Scientific Journal of Pharmacy)



JURNAL ILMIAH FARMASI
(SCIENTIFIC JOURNAL OF PHARMACY)

PIMPINAN UMUM/ PENANGGUNG JAWAB
Dekan Fakultas Matematika dan Ilmu Pengetahuan Alam
Universitas Islam Indonesia

WAKIL PIMPINAN UMUM/ WAKIL PENANGGUNG JAWAB
Ketua Jurusan Farmasi FMIPA UII

MITRA BESTARI

1. Prof. Dr. Wiryatun Lestariana, Apt
2. Prof. Dr. Zullies Ikawati, Apt
3. Prof. Dr. Sudibyo Martono, Apt
4. Dr. Tedjo Yuwono, Apt
5. Prof. Dr. Dachriyanus, Apt
6. Prof. dr. Iwan Dwiprahasto, MMedSc, PhD
7. Prof. Dr. Lukman Hakim M.Sc., Apt
8. Prof. Dr. Achmad Fudholi, DEA, Apt
9. Prof. Dr. Ibnu Gholib Gandjar, DEA., Apt

DEWAN EDITOR

Ketua : Saepudin, M.Si., Apt
Sekretaris : Rochmy Istikharah, M.Sc., Apt.
Anggota : Vitarani Dwi Ananda Ningrum, M.Si., Apt
Okti R. Mafruhah, MSc., Apt
Dimas Adhi Pradana, MSc., Apt.
Fithria DA. Suryanegara, MSc., Apt.
Ari Wibowo, S.Farm., Apt
Arba Pramudita Ramadani, MSc., Apt.
Oktavia Indrati, S.Farm., Apt.

Penerbit

Jurusan Farmasi Fakultas Matematika dan Ilmu Pengetahuan Alam
Universitas Islam Indonesia

Alamat Penerbit

Jurusan Farmasi FMIPA UII
Jl. Kaliurang Km. 14,4 Yogyakarta 55584
Telp. (0274) 896439 ext. 3047
Email: jif@uii.ac.id

DIAGNOSIS DINI TOKSOPLASMOSIS KONGENITAL DENGAN PCR CAIRAN AMNION

Umatul Khoiriyah

Bagian Parasitologi Fakultas Kedokteran Universitas Islam Indonesia

ABSTRACT

Toxoplasmosis is caused by infection with the protozoan *Toxoplasma gondii*. Congenital Toxoplasmosis is an infection resulting from the transplacental passage of the parasites from infected mother to the fetus. An acute infection in pregnant women associated with a range of outcomes from subclinical infection to intra uterine death. Early diagnosis is important to make decision, therapy or abortion. At present, prosedur diagnostic of congenital toxoplasmosis uses conventional methode especially serological test, but the conventional methode is time consuming, and has high risk. PCR amniotic fluid is one methode to make dignostic of conegenital toxoplasmosis. This is very sensitif and rapid (24 hours) and safer than serological tes of the fetalblood. PCR can be done from 15 week after gestation.

Key word: congenital toxoplasmosis- PCR- amniotic fluid.